

Bacterial Blight of Cotton During 2011 & 2012: Field Trash or Seed

Allen, T.W., Dodds, D., Golden, B.R., Lu,
S., Sciumbato, G.L., Wrather, A., Main,
C., Rothrock, C., Barber, T., Tzanetakis, I.,
and Kirkpatrick, T.

Cotton Incorporated,
Crop Management Seminar
Tunica, MS
November 8, 2012

Angular leaf spot/bacterial blight

- Caused by a bacterium (*Xanthomonas axonopodis* pv. *malvacearum*):
 - Leads to:
 - Seedling blight
 - Leaf spot
 - Vascular infection leads to blackened veins
 - Blackarm on stems and petioles
 - Boll blight (open wound for secondary organisms)
 - Boll rot (either the bacterium **OR** secondary)
- Foliar phase is commonly referred to as “angular leaf spot”
 - **BUT** not all angular spots are bacterial blight
 - Angular nature of the symptoms results from capillary veins limiting the movement of the organism











2011 disease distribution



2011 & 2012 disease distribution



2011 Arkansas Seed Testing

- Reported in: Rothrock et al., The Resurgence of Bacterial Blight on Cotton in Arkansas, 2012 Beltwide Cotton Conferences
- 2011 Arkansas infected varieties:
 - DP 0912, AM 1550, PHY 367, ST 5458
- 34 seed lots were screened from the above varieties
 - 14 of 34 seed lots contained *Xam* as an internal contaminant
 - 3 of 34 seed lots contained *Xam* as an external contaminant post-acid delinting

Bacterial blight facts

- Predominantly a seedborne organism
 - Reason for acid delinting
- ≈19-20 races of the bacterium worldwide
 - Virulence/pathogenicity differs between races
 - Mostly geographic in origin, the races in the U.S. are 1, 2, 5, 6, 7, 8, 11, 12, and 18 (18 is most common)
- Prefers high temperature (86-97°F) with high humidity (> 85%)
- Survival in soil depends on prevailing environment between seasons and tillage
- Yield loss (??%)


Variety	Rating (0-9)	Resistant (R) or Susceptible (S)
Americot 1550B2RF	4.7 bcd	S
DynaGro 2450B2RF	4.6 b-e	S
DynaGro 2570B2RF	4.6 b-e	S
Deltapine 0912B2RF	3.9 efg	S
Deltapine 1028B2RF	4.5 b-f	S
Deltapine 1034B2RF	4.7 bcd	S
Deltapine 1048B2RF	4.9 abc	S
Deltapine 1133B2RF	0.1 h	R
Deltapine 1137B2RF	5.4 a	S
Fibermax 1740B2F	0.2 h	R
Fibermax 9058F	0.2 h	R
HQ 210CT	4.2 d-g	S
HQ 212CT	3.8 fg	S
Phytogen 367WRF	4.8 a-d	S
Phytogen 375WRF	0.2 h	R
Phytogen 499WRF	5.2 ab	S
Phytogen 565WRF	4.4 c-f	S
Phytogen 725F	3.6 g	S
Stoneville 4145LLB2	3.8 fg	S
Stoneville 4288B2F	4.7 bcd	S
Stoneville 5288B2F	0.1 h	R
Stoneville 5458B2RF	4.5 c-f	S

Source: Allen, T.W., Golden, B., Sciumbato, G.L., and Thaxton, P. (unpublished) **NOTE:** $\alpha = 0.05$; $LSD=0.6793$; $CV=13.7$; $R^2=0.9553$

Impact on plant height



Yield (lint yield/Acre)



NOTE: $\alpha = 0.10$

2012 cotton field survey

Variety	# of fields with blight	# of fields considered	Percentage infected
DPL0912	66	88	75.0%
DPL0935	2	3	66.7%
DPL1028	28	28	100%
DPL1034	138	217	63.6%
DPL1048	28	37	75.7%
DPL1050	22	28	78.6%
PHY499	9	32	28.1%
ST5458	71	237	30.0%
TOTAL	364	670	54.3%

NOTE: information obtained from a survey of 13 observers in MS

Management Options

- Tillage
 - Bury stubble rather than minimum tillage practices
- Crop rotation (non-host)
 - Corn, grain sorghum, peanut, soybean
- Plant a resistant cotton variety
 - If the particular producer encountered bacterial blight in 2011 and does **NOT** want to change the tillage system, planting a resistant variety is the best option

Rapid Bacterial Blight Test Development

- Joint project between Univ. of Ar., MSU
- Presently, screening seed lots requires tremendous period of time (\approx 14-18 days)
- **Objectives:**
 - Rapid and accurate qPCR protocol
 - Field test the resulting methods
 - Comparisons with fungal leaf spots
 - Determine overall reaction of commercially available cotton varieties to bacterial blight pathogen



Tom Allen
DREC, Stoneville, MS
662-402-9995
tallen@drec.msstate.edu